



A Publication
of Reliable Methods
for the Preparation
of Organic Compounds

Working with Hazardous Chemicals

The procedures in *Organic Syntheses* are intended for use only by persons with proper training in experimental organic chemistry. All hazardous materials should be handled using the standard procedures for work with chemicals described in references such as "Prudent Practices in the Laboratory" (The National Academies Press, Washington, D.C., 2011; the full text can be accessed free of charge at http://www.nap.edu/catalog.php?record_id=12654). All chemical waste should be disposed of in accordance with local regulations. For general guidelines for the management of chemical waste, see Chapter 8 of Prudent Practices.

In some articles in *Organic Syntheses*, chemical-specific hazards are highlighted in red "Caution Notes" within a procedure. It is important to recognize that the absence of a caution note does not imply that no significant hazards are associated with the chemicals involved in that procedure. Prior to performing a reaction, a thorough risk assessment should be carried out that includes a review of the potential hazards associated with each chemical and experimental operation on the scale that is planned for the procedure. Guidelines for carrying out a risk assessment and for analyzing the hazards associated with chemicals can be found in Chapter 4 of Prudent Practices.

The procedures described in *Organic Syntheses* are provided as published and are conducted at one's own risk. *Organic Syntheses, Inc.*, its Editors, and its Board of Directors do not warrant or guarantee the safety of individuals using these procedures and hereby disclaim any liability for any injuries or damages claimed to have resulted from or related in any way to the procedures herein.


These paragraphs were added in September 2014. The statements above do not supersede any specific hazard caution notes and safety instructions included in the procedure.

Organic Syntheses, Coll. Vol. 1, p.149 (1941); Vol. 3, p.28 (1923).

CATECHOL

[Pyrocatechol]

[(A) (from Salicylaldehyde)]



Submitted by H. D. Dakin

Checked by H. T. Clarke and E. R. Taylor.

1. Procedure


To a solution of 122 g. (1 mole) of pure [salicylaldehyde](#) ([Note 1](#)) in 1000 cc. of normal [sodium hydroxide](#) solution at room temperature, is added 1420 g. (1.2 moles) of 3 per cent [hydrogen peroxide](#). The mixture darkens slightly in color and the temperature rises to 45–50°. The solution is allowed to stand for fifteen to twenty hours, whereupon a few drops of [acetic acid](#) are added in order to neutralize any excess alkali, and the solution evaporated to complete dryness on the water bath under reduced pressure.

The solid residue is finely crushed and warmed nearly to boiling with 500 cc. of [toluene](#); the mixture is then poured into the folded filter paper of an extraction apparatus ([p. 375](#)) and extracted with boiling [toluene](#) for five hours. The [toluene](#) is allowed to cool and is decanted from the [catechol](#), which crystallizes out. The insoluble material is again ground up and extracted in the apparatus with the decanted [toluene](#). The combined product, weighing 70–76 g., consists of light brown plates melting at 104°, and is thus pure enough for many purposes. A further 6–12 g. of [catechol](#) can be obtained on distilling off the bulk of the [toluene](#) from the mother liquor. In order to obtain an entirely pure product, the crude [catechol](#) should be distilled under reduced pressure, when it passes over entirely at 119–121°/10 mm. (or 113–115°/8 mm.), and the distillate recrystallized from about five times its weight of [toluene](#). In this way, colorless plates melting at 104–105° are obtained. The yield of the purified product is 76–80 g. (69–73 per cent of the theoretical amount) ([Note 2](#)).

2. Notes

1. A considerably lower yield is obtained (50 per cent or less) if technical [salicylaldehyde](#) (not purified through the bisulfite compound) is employed.
2. The procedure described is applicable to almost all hydroxyaldehydes in which the hydroxyl and carbonyl groups occupy ortho or para positions relatively to each other;¹ in the latter case derivatives of [hydroquinone](#) are produced. When the hydroxyl and carbonyl groups occupy the meta position with respect to each other, no reaction takes place, as is also the case with certain ortho and para compounds containing nitro groups and iodine atoms. [o-Hydroxyacetophenone](#) and [p-hydroxyacetophenone](#) are also capable of yielding [catechol](#) and [hydroquinone](#), respectively, under the above conditions.
3. [Catechol](#) may also be produced from [salicylaldehyde](#) by the use of certain derivatives of [hydrogen peroxide](#), such as persulfates or [sodium peroxide](#), but the method is far less convenient.

[(B) (from Guaiacol)]




Submitted by H. T. Clarke and E. R. Taylor.
Checked by C. S. Marvel and W. B. King.

1. Procedure

In a 5-l. flask fitted with a stirrer, a thermometer reaching to the bottom, a fractionating column about 40 cm. long (Note 1), and a return inlet tube, are placed 912 g. (7.4 moles) of crystalline [guaiacol](#) and 1500 g. (1015 cc., 8.8 moles) of 48 per cent [hydrobromic acid](#) (p. 26). This apparatus is connected with a condenser, an automatic separator, and an absorption vessel, as shown in Fig. 10 (Note 2). The mixture is gently heated, with continual stirring, at such a rate that the vapor at the head of the column maintains a temperature of 85–95°. The mixture of water and [guaiacol](#) which condenses passes through the automatic separator (Note 3), the [guaiacol](#) being returned to the reaction vessel and the water collected in the tap funnel, from which it is removed from time to time. The gaseous [methyl bromide](#) is collected in an ice-cooled bottle containing a weighed quantity (about 1600 g.) of [methyl alcohol](#) (Note 4).

Fig. 10.



After six to seven hours, the temperature at the head of the column begins to rise above 95° and finally reaches 98°, at which point practically no more [guaiacol](#) passes over with the distillate. This requires about one hour more, after which the heating is stopped (Note 5). The increase in weight of the absorption bottle amounts to 400–500 g. (57–72 per cent of the theoretical amount of [methyl bromide](#)); the water collected weighs 550–560 g. (Note 6).

The residue in the flask, which has become pink in color, is extracted at 85–95° with three 1500-cc. portions of [toluene](#) (Note 7), and the united extracts are distilled under reduced pressure on the steam bath until no more moisture passes over with the vapor. The solution (2–2.5 l.) is then allowed to cool,

whereupon 385–390 g. of **catechol** separates in colorless plates melting at 104–105°. The mother liquor is further concentrated under reduced pressure and the residue finally distilled, pure **catechol** passing over at 124–125° / 12 mm. The distillate, on recrystallization from a small quantity of **toluene**, yields 8–10 g. of pure material.

The aqueous residue from the three extractions with **toluene** yields, when the bulk of the water has been distilled off on the steam bath under reduced pressure, a further quantity of **catechol** by extraction with hot **toluene**, but the crystals so obtained are contaminated with a red impurity. It is therefore preferable to distil the entire residue, again collecting the fraction boiling at 124–125° / 12 mm. and recrystallizing it from fresh **toluene**. In this way 295–305 g. of pure product is obtained. The forerun contains a little aqueous **hydrobromic acid**, which may be employed in a subsequent batch. The total yield of pure **catechol** is 690–705 g. (85–87 per cent of the theoretical amount) (Note 8).

2. Notes

1. Any column of standard form is suitable; to work satisfactorily, it is unnecessary for it to be more than 40 cm. in length.
2. In the diagram (Fig. 10) no indication is made of joints in the glass tubing. These are, however, applicable at almost any desired point; but care should be taken to arrange them so that the ends of the glass tube come into contact, since **methyl bromide** appears to pass readily through the walls of the rubber tubing. It is necessary to make certain that the outlets from the automatic separator descend in an even incline and that the tubing itself is of sufficiently large diameter to prevent any siphoning effects.
3. It is noteworthy that the **guaiaicol** which condenses is saturated with **methyl bromide** at the condensation temperature, so that it is necessary to prevent it from becoming warmer during its passage through the separator; otherwise the evolution of gas may hinder the ready formation of layers.
4. In order to secure satisfactory absorption of the **methyl bromide**, the **methyl alcohol** should be kept below 5° by ice-cooling.
5. Since the formation of the red by-product appears to depend upon the time during which the **catechol** is heated with **hydrobromic acid**, it is well to carry out the reaction as rapidly as possible; due care must be taken, however, that the temperature at the head of the column does not rise prematurely above 95°.
6. The water collected during the reaction contains practically no **hydrobromic acid** but is saturated with **guaiaicol** and **methyl bromide**. Nevertheless, it scarcely pays to recover the dissolved compounds, for on distillation only 1–2 g. of wet **guaiaicol** can be collected; while the value of the **methyl bromide** is in any case not very great.
7. It is suggested that **carbon tetrachloride** rather than **toluene** be used in the extraction because **toluene** gives a product which gradually colors, whereas the **catechol** extracted with **carbon tetrachloride** remains colorless for a considerably longer time (W. W. Hartman, private communication).
8. The reaction appears to be applicable to many other cases where the constitution of the molecule permits of boiling with strong **hydrobromic acid**.

3. Discussion

Catechol can be prepared by the oxidation of **salicylaldehyde**¹ and of **phenol**;² by the demethylation of **guaiaicol** with **aluminum chloride**³ and **hydriodic acid**;⁴ by hydrolysis of ortho halogen-substituted phenols or ortho dihalobenzenes under suitable conditions;⁵ and by the replacement of sulfonic groups by alkali fusion.⁶ A series of reactions involving the oxidation of **safrole**, opening of the methylenedioxy ring, and decarboxylation also furnishes **catechol**.⁷

This preparation is referenced from:

- [Org. Syn. Coll. Vol. 3, 759](#)

References and Notes

1. Dakin, *Am. Chem. J.* **42**, 477 (1909).

2. Martinon, Bull. soc. chim. (2) **43**, 157 (1885); Magidson, Porozovska and Seligsohn, Trans. Sci. Chem. Pharm. Inst. Moscow, **6**, 23 (1923) [C. A. **22**, 3884 (1928)].
 3. Hartmann and Gattermann, Ber. **25**, 3532 (1892); Bayer and Co., Ger. pat. 70,718 [Frđl. **3**, 52 (1890–94)].
 4. Baeyer, Ber. **8**, 153 (1875); Perkin, J. Chem. Soc. **57**, 587 (1890).
 5. Lautemann, Ann. **120**, 315 (1861); Körner, Z. Chem. **322** (1868); Merck, Ger. pat. 84,828 [Frđl. **4**, 114 (1894–97)]; Friedr. Bayer and Co., Ger. pat. 249,939 [Frđl. **10**, 1330 (1910–12)]; Boehringer and Sons, Ger. pat. 269,544 [Frđl. **11**, 190 (1912–14)]; Fittig and Mager, Ber. **8**, 363 (1875); Kipriyanov and Suich, Ukrain. Khem. Zhur. **7**, Wiss.-tech. Abt. 94 (1932) [C. A. **27**, 3824 (1933)]; E. I. du Pont de Nemours and Co., Brit. pat. 425,230 [C. A. **29**, 5458 (1935)].
 6. Kekulé, Z. Chem. **643** (1867); Merck, Ger. pat. 80,817 [Frđl. **4**, 116 (1894–97)]; Tobias, Ger. pat. 81,209 [Frđl. **4**, 117 (1894–97)].
 7. Hiraizumi, J. Chem. Soc. Japan **55**, 601 (1934) [C. A. **28**, 5422 (1934)].
-

Appendix
Chemical Abstracts Nomenclature (Collective Index Number);
(Registry Number)

ortho halogen-substituted phenols

ortho dihalobenzenes

acetic acid (64-19-7)

methyl alcohol (67-56-1)

sodium hydroxide (1310-73-2)

hydroquinone (123-31-9)

phenol (108-95-2)

HYDROBROMIC ACID (10035-10-6)

carbon tetrachloride (56-23-5)

aluminum chloride (3495-54-3)

toluene (108-88-3)

Catechol,
Pyrocatechol (120-80-9)

Salicylaldehyde (90-02-8)

hydrogen peroxide (7722-84-1)

sodium peroxide

Guaiacol (90-05-1)

methyl bromide (74-83-9)

hydriodic acid (10034-85-2)

safrole (94-59-7)

o-Hydroxyacetophenone (118-93-4)

p-hydroxyacetophenone (99-93-4)