



A Publication
of Reliable Methods
for the Preparation
of Organic Compounds

Working with Hazardous Chemicals

The procedures in *Organic Syntheses* are intended for use only by persons with proper training in experimental organic chemistry. All hazardous materials should be handled using the standard procedures for work with chemicals described in references such as "Prudent Practices in the Laboratory" (The National Academies Press, Washington, D.C., 2011; the full text can be accessed free of charge at http://www.nap.edu/catalog.php?record_id=12654). All chemical waste should be disposed of in accordance with local regulations. For general guidelines for the management of chemical waste, see Chapter 8 of Prudent Practices.

In some articles in *Organic Syntheses*, chemical-specific hazards are highlighted in red "Caution Notes" within a procedure. It is important to recognize that the absence of a caution note does not imply that no significant hazards are associated with the chemicals involved in that procedure. Prior to performing a reaction, a thorough risk assessment should be carried out that includes a review of the potential hazards associated with each chemical and experimental operation on the scale that is planned for the procedure. Guidelines for carrying out a risk assessment and for analyzing the hazards associated with chemicals can be found in Chapter 4 of Prudent Practices.


The procedures described in *Organic Syntheses* are provided as published and are conducted at one's own risk. *Organic Syntheses, Inc.*, its Editors, and its Board of Directors do not warrant or guarantee the safety of individuals using these procedures and hereby disclaim any liability for any injuries or damages claimed to have resulted from or related in any way to the procedures herein.

These paragraphs were added in September 2014. The statements above do not supersede any specific hazard caution notes and safety instructions included in the procedure.

Organic Syntheses, Coll. Vol. 7, p.276 (1990); Vol. 63, p.203 (1985).

ONE-STEP HOMOLOGATION OF ACETYLENES TO ALLENES: 4-HYDROXYNONA-1,2-DIENE

[1,2-Nonadien-4-ol]



Submitted by Pierre Crabbé, Bahman Nassim, and Maria-Teresa Robert-Lopes¹.
Checked by Jeffrey S. Stults and Edwin Veddejs.

1. Procedure

In a 500-mL, three-necked flask, equipped with a thermometer, stirrer, and a reflux condenser with drying tube, are placed 12.6 g (0.1 mol) of **1-octyn-3-ol**, 154 mL of **dioxane**, 7.24 g (0.0504 mol) of **cuprous bromide**, 7.4 g of **paraformaldehyde**, and 18.54 g (0.183 mol) of **diisopropylamine** (Note 1). The resulting mixture is gently refluxed and stirred for 2 hr and then cooled to room temperature and filtered through a Celite plug. The dark-brown filtrate is concentrated under vacuum (Rotavapor) to a gummy residue and then diluted with 50 mL of water followed by 100 mL of **ether** and acidified with 6 *N* **hydrochloric acid** to pH 2. The ether–water layers are decanted from any residue, the **ether** layer is separated, and the aqueous solution is extracted with **ether** (5 × 50 mL). The **ether** extracts are combined and washed with small portions of water until pH 6.5 is reached. The organic layer is then washed with saturated **sodium chloride** solution and dried over anhydrous **MgSO₄**. After removal of **ether** by distillation through a 20-cm Vigreux column (water aspirator vacuum) while heating on a water bath, ≤ 40°C, the residual liquid is fractionated under reduced pressure through a 10-cm Vigreux column. The main fraction is collected at 41–42.5°C(0.15 mm) to give 8.65 g of pure allene (Note 2), with additional fractions of a less pure material.

2. Notes

1. **Cuprous bromide** and **1-octyn-3-ol** were used as supplied by the Aldrich Chemical Company, Inc. **Dioxane** was dried over **sodium–benzophenone** and distilled, and **diisopropylamine** was distilled from **barium oxide**.
2. The spectral properties of **4-hydroxynona-1,2-diene** are as follows: IR (neat) cm^{-1} : 3500 (OH), 1960 (C=C=C), 850 (=CH), 2900–2850 (CH). ¹H NMR (CDCl₃) δ : 0.65–1.7 (m); 4.15 (1 H, m); 4.8 (2 H, d of d, *J* = 2.6 Hz); 5.22, (1 H, q, *J* = 6 Hz).

3. Discussion

Although allenes were characterized long ago as a distinct class of organic substances, they have only recently received proper attention from chemists, in particular for their potential in organic synthesis.^{2–4} A number of methods are known for the transformation of acetylenes into allenes,⁵ but few are known to allow the homologation of an acetylenic group into a **propadiene** functionality.

A general procedure for the homologation of acetylenic compounds into allenes is described. The reaction conditions are mild and appear to be general, so that they can be applied to plain acetylenic substances as well as to acetylenic alcohols, ethers, and esters. This procedure is essentially a one-step reaction. As such, it is simpler and faster than the previously reported technique that involves the conversion of an acetylenic compound into the Mannich base, the formation of its quaternary ammonium salt and the reduction of this salt with **lithium aluminum hydride**.⁶ Of great advantage over previously available methodology are the mild conditions, as well as the clean and fast procedure, which make this a method of choice for an efficient conversion of acetylenes to allenes.^{7–9}

References and Notes

1. Department of Chemistry, University of Missouri, Columbia, MO 65211.
 2. Taylor, D. R. *Chem. Rev.* **1967**, *67*, 317 and references cited therein.
 3. Griesbaum, K. *Angew. Chem. Int. Ed. Engl.* **1966**, *5*, 933;
 4. Mavrov, M. V.; Kucherov, V. F. *Russ. Chem. Rev.* **1967**, *36*, 233.
 5. Fischer, H. In "The Chemistry of Alkenes," Patai, S., Ed.; Wiley-Interscience: New York, 1964; Chapter 13, p. 1025.
 6. Galantay, E.; Basco, I.; Coombs, R. V. *Synthesis* **1974**, 344.
 7. Crabbé, P.; André, D.; Fillion, H. *Tetrahedron Lett.* **1979**, 893;
 8. Crabbé, P.; Fillion, H.; André, D.; Luche, J. L. *J. Chem. Soc., Chem. Commun.* **1979**, 859;
 9. Searles, S.; Li, Y.; Nassim, B.; Lopes, M. T.; Tran, P. T.; Crabbé, P. *J. Chem. Soc., Perkin Trans. I* **1984**, 747.
-

Appendix Chemical Abstracts Nomenclature (Collective Index Number); (Registry Number)

hydrochloric acid (7647-01-0)

ether (60-29-7)

sodium chloride (7647-14-5)

barium oxide

Benzophenone (119-61-9)

sodium (13966-32-0)

cuprous bromide (7787-70-4)

MgSO₄ (7487-88-9)

dioxane (5703-46-8)

propadiene (463-49-0)

lithium aluminum hydride (16853-85-3)

diisopropylamine (108-18-9)

4-Hydroxynona-1,2-diene,
1,2-Nonadien-4-ol (73229-28-4)

1-octyn-3-ol (818-72-4)

paraformaldehyde (30525-89-4)

